

Global Health Challenges and Sustainable Healthcare System Development

Ryunosuke Irie and Koji Mikami*

Department of Integrative Studies of Plant and Animal Production, School of Food Industrial Sciences, Miyagi University, 2-2-1 Hatatate, Taihaku-ku, Sendai 982-0215, Japan

Abstract

The conchocelis, the filamentous sporophyte generation of Bangiales, proliferates through tip growth in which only the tips of branch initial cells elongate and subsequently divide perpendicularly to the apical–basal axis to form filamentous structures. As was found in terrestrial plants, the tip growth of conchocelis in the marine red alga *Pyropia yezoensis* is regulated by auxin, actin polymerization, and phosphatidylinositol signaling. Although extracellular Ca^{2+} influx is critical for tip growth in terrestrial plants, little is known about its involvement in the tip growth of algae. We therefore investigated whether extracellular Ca^{2+} influx is required for tip growth in *P. yezoensis* conchocelis. Treatment of isolated single-celled conchocelis with a Ca^{2+} chelator, ethylene glycol tetraacetic acid, inhibited both the formation of branch initials and the tip growth of branches in a dose-dependent manner. These findings indicate that extracellular Ca^{2+} influx is indeed involved in the tip growth of conchocelis, suggesting that the basic regulatory mechanisms governing tip growth may be conserved between marine red algae and terrestrial plants. Further confirmation of this possibility will require the characterization of the spatiotemporal patterns of Ca^{2+} oscillations and F-actin accumulation, as well as the subcellular localization of phosphoinositides and their catabolic enzymes in the *P. yezoensis* conchocelis.

Keywords: Branch initial, Ca^{2+} , Conchocelis, Extracellular influx, *Pyropia yezoensis*, Red alga, Tip growth

Corresponding author:

Koji Mikami, Department of Integrative Studies of Plant and Animal Production, School of Food Industrial Sciences, Miyagi University, 2-2-1 Hatatate, Taihaku-ku, Sendai 982-0215, Japan, E-mail: mikamik@myu.ac.jp

Received: September 02, 2022; **Accepted:** September 19, 2022; **Published:** September 27, 2022

Citation: Canam T, et al. Navigating the Global Nursing Shortage: Implications for Basic Care Nursing Services. Research Chronicle in Health Sciences. 2022, Vol. 8 No. 2: 105

Introduction

Patterns of anisotropic cell growth give rise to the body shapes of terrestrial plants and multicellular algae [1,2]. Diffuse growth, which results in the expansion of cell surfaces [3,4], produces planar or foliose shape, whereas tip growth [5-7], characterized by directional and highly localized expansion at the apex, produces a filamentous body architecture. Algal cells grow anisotropically through both diffuse and tip growth. We previously demonstrated that tip growth produces the filamentous structures of the conchocelis (sporophyte) and conchosporangium (conchosporophyte) of the red alga *Pyropia yezoensis* [8-12], although regulatory mechanisms of tip growth in *P. yezoensis* has not yet been fully elucidated. Moreover, how diffuse growth produces a foliose shape in the thallus (gametophyte) of *P.*

yezoensis remains largely unknown. Therefore, characterizing the contrasting mechanisms that regulate patterns of diffuse and tip growth is necessary to understand the differences in growth–morphology relationships between the two life-cycle generations of Bangiales.

The mechanisms that regulate diffuse and tip growth systems have been studied extensively in terrestrial plants. Cortical microtubules (MTs), actin filament (F-actin), Rho GTPases, the histone variant H2A.Z, gibberellin, and brassinosteroids have all been implicated in the regulation of diffuse growth [1,3,13] (Likewise, MTs, F-actin, Rho GTPases, Ca^{2+} influx, reactive oxygen species, phosphatidylinositol signaling, auxin, and jasmonate have been shown to regulate tip growth [7,13-21]. Although research on the regulation of diffuse growth in macroalgae

has shown little progress, recent studies have provided basic information about the mechanisms regulating tip growth in red and brown macroalgae [2,11,12,22]. Indeed, we demonstrated that auxin, phosphoinositide turnover, and actin polymerization are involved in the regulation of tip growth in *P. yezoensis* [11,12], consistent with findings in terrestrial plants [13,17,18,21]. Although extracellular Ca^{2+} influx is required for tip growth in terrestrial plants [17-19], little is known about its role in the tip growth of macroalgae. We therefore sought to determine whether extracellular Ca^{2+} influx is required for tip growth in *P. yezoensis* conchocelis.

Materials and Methods

Single-celled conchocelis of *P. yezoensis* strain U-51 were used for these experiments, enabling us to visualize and quantitatively analyze the initiation and progression of tip growth in detail [11]. Conchocelis were maintained in artificial seawater at 15°C under 60 $\mu\text{E}/\text{m}^2/\text{s}$ of light with a short-day cycle (10-h light/14-h dark) [23]. Single-celled conchocelis were prepared as described in [11]. In brief, aggregates of multicellular conchocelis were chopped with a razor blade, and small fragments of conchocelis were separated from larger pieces by filtration through a 10- μm nylon mesh. The separated conchocelis were then incubated in 30 mL of artificial seawater [23] at 15°C for 10 min. Unbranched single-celled conchocelis (Figure 1) were identified by observation under an Olympus IX73 light microscope equipped with an Olympus DP22 camera (Olympus Corporation, Tokyo, Japan), drawn into a micropipette, and transferred to 96-well plates (one cell per well containing 200 μL of artificial seawater). Ethylene glycol tetraacetic acid (EGTA; Dojindo Laboratories, Japan), an effective Ca^{2+} chelator, was dissolved in artificial seawater to

create a 0.5 M stock solution (adjusted to pH 8.0 with NaOH) and stored at -30°C before use. Single-celled conchocelis were treated with 0, 500, or 750 μM EGTA for 3 days, and the growth and morphology of the side branches were observed using the microscope described above.

Results and Discussion

To evaluate the effects of EGTA on the initiation and elongation of side branches, we measured the branching rate (the percentage of observed cells that produced side branches) and branch length, respectively. Side branches were initiated and underwent elongation when treated with 0 (control) or 500 μM EGTA (Figure 2A and 2B), but 750 μM EGTA completely prevented both the initiation and growth of side branches (Figure 2C). The branching rate and branch length were affected by the presence of EGTA in a dose-dependent manner, being strongly inhibited at 750 μM (Figure 2D and 2E). These results demonstrate that extracellular Ca^{2+} influx is important for the production and tip growth of branch initials from differentiated conchocelis cells. It reminds to be elucidated whether washing out of EGTA and subsequent addition of CaCl_2 in the medium recover branch initiation and branch growth and whether inhibition of channel-mediated Ca^{2+} influx by treatment with LaCl_3 prevents tip growth.

There are two important steps in the tip growth of conchocelis: the formation of branch initials in non-dividing differentiated cells (initiation of tip growth) and the polar directional growth of these initials to form elongated branches [12]. Because the treatment of single-celled conchocelis with EGTA impaired both branch initiation and branch growth, we concluded that Ca^{2+} is required for both critical steps in tip growth. To confirm this proposal, measurements of the time-course of both production of branch initial and its growth in single-celled conchocelis are necessary for elucidation of a question which step requires extracellular Ca^{2+} influx. In addition, we previously demonstrated that these two steps are regulated by auxin, phosphoinositide turnover, and actin polymerization [11,12]. These findings are consistent with our understanding of tip-growth regulation in the pollen tubes and root hairs of terrestrial plants [13,14,16-18,21]. It is therefore possible that the basic regulatory mechanisms governing tip growth are conserved between aquatic red algae and terrestrial plants. Factors such as Ca^{2+} , enzymes related to phosphoinositide turnover, and F-actin act at the cell apex to enable the polar growth of filamentous tissues such as root hairs and pollen tubes [14,16,18]; however, little is known about the spatial distribution of these factors during the initiation and elongation of branches in *P. yezoensis* conchocelis. Therefore, determining the specific regions where auxin and Ca^{2+} exert their effects and characterizing the subcellular distributions of F-actin and enzymes involved in phosphoinositide turnover will be critical for fully understanding tip-growth regulation in conchocelis branches. A live-imaging technique for conchocelis tip growth has recently been reported




Figure 1: Representative image of an isolated single-celled conchocelis. Nonbranched single-celled conchocelis were used for EGTA treatment experiments. Scale bar, 25 μm .




Figure 2: Effects of EGTA treatment on the tip growth of side branches from single-celled conchocelis. (A–C) Photographs of single-celled conchocelis treated with 0 (A), 500 (B), or 750 μM EGTA (C) for 3 days, for which each treatment was employed total 16 isolated

[10]. Improvements to this system are expected to enable the visualization of Ca^{2+} influx, F-actin distribution, and the localization of phosphoinositides and related enzymes, providing spatiotemporal details of polarity establishment and tip growth in *P. yezoensis* conchocelis.

Despite progress in understanding tip growth in the conchocelis, much less is known about the regulation of diffuse growth that gives rise to the foliose thallus of the gametophyte. It is remarkable that foliose and filamentous generations survive independently in *P. yezoensis* and other Bangiales [9] (suggesting differences in the regulatory mechanisms governing these two growth patterns. However, the shared involvement of MTs, F-actin, and Rho GTPases in both diffuse and tip growth of terrestrial plants [1] raises the question of how common factors can regulate different types of growth. Characterizing the contrasting regulatory mechanisms that govern diffuse and tip growth will be essential for understanding the physiological and molecular bases of the distinct multicellular body plans of heteromorphic generations in *P. yezoensis*.

Author Contributions

RI: Methodology, Investigation, Data curation, Formal analysis, Validation, Visualization.

KM: Conceptualization, Methodology, Formal analysis, Validation, Visualization, Supervision, Writing—original draft preparation and reviewing and editing, Funding acquisition.

Conflict of Interest

The authors declare no conflicts of interest. No aspects of the study required informed consent, and the work did not involve human or animal subjects. All authors have read and agreed to authorship and to the submission of the manuscript for peer review.

Data Availability Statement

Data is contained within the article.

References

- Guerriero G, Hausman JF, Cai G (2014) No stress! Relax! Mechanisms governing growth and shape in plant cells. *Int J Mol Sci* 15(3): 5094-114.
- Rabillé H, Billoud B, Tesson B, Le Panse S, Rolland É, et al. (2019) The brown algal mode of tip growth: Keeping stress under control. *PLoS Biol* 17(1): e2005258.
- Braidwood L, Breuer C, Sugimoto K (2014) My body is a cage:

- mechanisms and modulation of plant cell growth. *New Phytol* 201(2): 388-402.
4. Cosgrove DJ (2018) Diffuse growth of plant cell walls. *Plant Physiol* 176(1): 16-27.
 5. Dumais J, Shaw SL, Steele CR, Long SR, Ray PM (2006) An anisotropic-viscoplastic model of plant cell morphogenesis by tip growth. *Int J Dev Biol* 50(2-3): 209-222.
 6. Rounds CM, Bezanilla M (2013) Growth mechanisms in tip-growing plant cells. *Annu Rev Plant Biol* 64: 243-265.
 7. Bibeau JP, Galotto G, Wu M, Tüzel E, Vidali L (2021) Quantitative cell biology of tip growth in moss. *Plant Mol Biol* 107(4-5): 227-244.
 8. Takahashi M, Mikami K (2016) Phototropism in the marine red macroalga *Pyropia yezoensis*. *Am J Plant Sci* 7: 2412-2428. <https://doi.org/10.4236/ajps.2016.717211>
 9. Mikami K, Li C, Irie R, Hama Y (2019) A unique life cycle transition in the red seaweed *Pyropia yezoensis* depends on apoptosis. *Commun Biol* 2: 299.
 10. Hiwatashi Y, Shimada M, Mikami K, Takada N (2022) Establishment of a live-imaging analysis for polarized growth of conchocelis in the multicellular red alga *Neopyropia yezoensis* *Front Plant Sci* 12: 716011.
 11. Taya K, Takeuchi S, Takahashi M, Hayashi K, Mikami K (2022) Auxin regulates apical stem cell regeneration and tip growth in the marine red alga *Neopyropia yezoensis*. ***Cells* 11(17): 2652.**
 12. Irie R, Mikami K (2025) Phosphoinositide signaling and actin polymerization are critical for tip growth in the marine red alga *Pyropia yezoensis*. *Plants* **14(14): 2194.**
 13. Ma C, Chang J, Hazelwood OS, Arif Ashraf M, Nan Q (2020) Establishing cell polarity in plants: the role of cytoskeletal structures and regulatory pathways. *Front Cell Dev Biol* 13: 1602463.
 14. Ischebeck T, Seiler S, Heilmann I (2010) At the poles across kingdoms: phosphoinositides and polar tip growth. *Protoplasma*; 240(1-4): 13-31.
 15. Saavedra L, Mikami K, Malhó R, Sommarin M (2012) PIP kinases and their role in plant tip growing cells. *Plant Signal Behav* 7(10): 1302-1305.
 16. Qu X, Jiang Y, Chang M, Liu X, Zhang R, Huang S (2015) Organization and regulation of the actin cytoskeleton in the pollen tube. *Front Plant Sci* 5: 786.
 17. Lee YJ, Yang Z (2008) Tip growth: signaling in the apical dome. *Curr Opin Plant Biol* 11(6): 662-671.
 18. Scholz P, Anstatt J, Krawczyk HE, Ischebeck T (2020) Signalling pinpointed to the tip: the complex regulatory network that allows pollen tube growth. *Plants* 9: 1098.
 19. Zhang X, Bian A, Li T, Ren L, Li L, Su Y, Zhang Q (2022) ROS and calcium oscillations are required for polarized root hair growth. *Plant Signal Behav* 17(1): 2106410.
 20. Han X, Kui M, He K, Yang M, Du J, et al. (2020) Jasmonate-regulated root growth inhibition and root hair elongation. *J Exp Bot* 74: 1176-1185.
 21. Berdion Gabarain V, Núñez-Lillo G, Pěnčík A, Carignani Sardoy M, Ibeas MA, et al. (2020) Enhanced auxin signaling promotes root-hair growth at moderately low temperature in *Arabidopsis thaliana*. *Plant Commun* 6(6): 101350.
 22. Charrier B, Le Bail A, de Reviers B (2012) Plant Proteus: brown algal morphological plasticity and underlying developmental mechanisms. *Trends Plant Sci* 17(8): 468-77. []
 23. Li C, Ariga I, Mikami K (2019) Difference in nitrogen starvation-inducible expression patterns among phylogenetically diverse ammonium transporter genes in the red seaweed *Pyropia yezoensis*. *Am J Plant Sci*, 10: 1325–1349. <https://doi: 10.4236/ajps.2019.108096>